

GLOBAL CONSTRAINTS OF GAUSS' LAW AND THE SOLUTION TO THE STRONG CP PROBLEM

HUAZHONG ZHANG

*P.O.Box 17660, Jackson State University
Jackson, MS 39217, USA*

ABSTRACT

This is a brief review on the work done recently. It is shown that the global constraints of Gauss' law ensure that the vacuum angle must be quantized in gauge theories with magnetic monopoles. Our quantization rule is given as $\theta = 0$, or $\theta = 2\pi N/n$ ($n \neq 0$) with integer n being the relevant topological charge of the magnetic monopole and N is an unfixed integer in this approach. This theoretically confirms further the conclusion originally proposed by the author¹ that the strong CP problem can be solved due to the existence of magnetic monopoles, the fact that the strong CP-violation can be only so small or vanishing may imply the existence of magnetic monopoles, and the universe is open.

1. Introduction and the Summary of the Main Results

It is well known that the effective θ term in QCD with an arbitrary effective and chiral invariant vacuum angle $\theta + \arg(\det M_{quark})$, which will be denoted as θ hereafter can induce CP violations in strong interactions. However, the possible electric dipole moment of the neutron puts a limit $\theta \leq 10^{-9}$ (modulo 2π). The need for understanding this leads to the strong CP problem. Recently, a non-perturbative solution to the Strong CP problem with magnetic monopoles has been proposed originally by the author^{1,2}. This may be interesting since the axions needed in the well-known Peccei-Quinn solution with an additional U(1) symmetry³ have not been observed. In our solution^{1,2}, it is proposed that the vacuum angle with magnetic monopoles must be quantized. Our quantization rule is derived by two different methods. This is given by $\theta = 0$, or $\theta = 2\pi N/n$ ($n \neq 0$) with integer n being the relevant topological charge of the magnetic monopole and N is an integer. In Ref. 1, it is shown that the integer N can be chosen as 1. The first method¹ is to show the existence of a monopole structure in the relevant gauge orbit space in Schrodinger formulation^{4,5}, and using the Dirac quantization rule for having a well-defined wave functional. The relevance to the $U_A(1)$ problem and the relevant vortex structures are also discussed in Ref.1. The second method is to show that there exist well-defined gauge transformations which will ensure the quantization of θ by the constraints of Gauss's law due to the non-abelian electric charges carried by the magnetic monopoles proportional to θ as noted in Ref. 6 and generalized in Ref. 7.

Therefore, we conclude that strong CP problem can be solved due to the quantization of θ in the presence of magnetic monopoles, for example monopoles of topological charge $n = \pm 1$ with $\theta = \pm 2\pi$, or $n \geq 2\pi 10^9$ with $\theta \leq 10^{-9}$. Moreover, the existence of non-vanishing magnetic flux through the space boundary implies that the universe must be open. In this note, we will briefly discuss and review the method in Ref. 2.

2. Constraints of Gauss' Law and the Qantization Rule for θ

In the Schrodinger formulation, the gauge theory system is described by a Hamiltonian equation with the constraints of Gauss' law for the wave functional $\Psi[A]$. Consider the non-abelian gauge theories with a θ term with a non-singular magnetic monopole at the origin. Such a θ term in the presence of magnetic monopole was first considered⁸ relevant to $U_A(1)$ problem. It was noted by Witten and generalized in Ref. 7 that monopoles of magnetic charges $q = G_a L_a$, can carry induced non-abelian electric charges $q_e = m \frac{q}{2\pi} - \frac{\theta q}{4\pi^2}$, $m \in Z$, where L_a ($a=1, 2, \dots \dim(G)$) form a basis for the Lie algebra of the gauge group. For an arbitrary value of θ , monopoles must be dyons. For simplicity, we will consider the case of static field for our purpose. The θ term can be written as

$$\mathcal{L}_\theta = \frac{\theta}{4\pi^2} \mathbf{E} \cdot \mathbf{B}. \quad (1)$$

Where $\mathbf{E}^a = -\nabla \phi^a - C_{abc} \mathbf{A}^b \phi^c$, $\mathbf{B}^a = \nabla \times \mathbf{A}^a - \frac{1}{2} C_{abc} \mathbf{A}^b \times \mathbf{A}^c$ with $\phi^a = A_0^a$, and C_{abc} being the structure constants of the Lie algebra. With integration by part, the θ term induces a non-abelian electric charge density

$$\rho^a = \frac{\theta}{4\pi^2} \nabla \cdot \mathbf{B}^a - \frac{\theta}{4\pi^2} C_{abc} \mathbf{A}^b \cdot \mathbf{B}^c. \quad (2)$$

The first and the second terms will induce a non-abelian electric charge at the monopole^{6,7} and a non-abelian electric charge density in the space outside the monopole respectively⁷.

We will choose the Weyl gauge $A_0 = 0$. The constraints of Gauss' law for $\Psi_{phys}[\mathbf{A}]$ is given by

$$\{-i D_i^{ab} \frac{\delta}{\delta A_i^b} - \rho^a\} \Psi_{phys}[\mathbf{A}] = 0, \quad (3)$$

where $D_i^{ab} = \delta^{ab} \partial_i + C_{abc} A_i^c$. Now multiply the above equation for the constraints of Gauss law by infinitesimal $\lambda^a(\mathbf{x})$, sum over a and integrate over $\mathbf{x}$. Then we obtain

$$\int d^3x \{i (D_i \lambda)^a \frac{\delta}{\delta A_i^a} - \lambda^a \rho^a\} \Psi_{phys}[\mathbf{A}] = 0. \quad (4)$$

This gives $\delta_\lambda \Psi_{phys}[\mathbf{A}] = \{-i \int d^3x \lambda^a \rho^a\} \Psi_{phys}[\mathbf{A}]$. For a well-defined finite gauge transformation with gauge function $g(\mathbf{x}) = \exp\{\lambda^a(\mathbf{x}) L_a\}$, we can consider the path $g(\mathbf{x}, t) = \exp\{t \lambda^a(\mathbf{x}) L_a\}$ $t \in [0, 1]$, connecting $g(\mathbf{x})$ and the identity element e . Consider a closed

loop C^1 with $g(\mathbf{x}, 1) = g(\mathbf{x}, 0) = e$, the state wave functional $\Psi_{phys}[\mathbf{A}]$ is invariant. Then by integration along the closed path⁴, with $\mathbf{A}^g = g^{-1}\mathbf{A}g + g^{-1}\nabla g$ and $tr(L_a L_b) = -\frac{1}{2}\delta_{ab}$, in terms of field strength two form F , this gives² topologically

$$\frac{\theta}{2\pi^2} tr \left\{ \int_{S^2} F \int_{C^1} \delta g g^{-1} \right\} = 2\pi k, \quad (5)$$

with k being integers. Now the magnetic charges can be transformed into a given Cartan subalgebra with the basis H_i ($i=1, 2, \dots, r$) as

$$\int_{S^2} F = G_0 = 4\pi \sum_{i=1}^r \beta^i H_i. \quad (6)$$

The constraints from the well-defined gauge transformations must be physical. Consider the following⁹ well-defined $U(1)$ gauge subgroup on the space boundary given by

$$g(t) = \exp \left\{ 4\pi m t \sum_{i,j} \frac{(\alpha_i)^j H_j}{\langle \alpha_i, \alpha_i \rangle} \right\}, \quad m \in Z, \quad t \in [0, 1], \quad (7)$$

where the α_i ($i=1, 2, \dots, r$) are the simple roots of the Lie algebra. We obtain $\exp \left\{ i \frac{\theta n m}{2\pi} \right\} = 1$. Where n is the effective topological number of the magnetic monopole defined as¹

$$n = -2 \langle \delta', \beta \rangle = -2 \sum_{i=1}^r \frac{\langle \alpha_i, \beta \rangle}{\langle \alpha_i, \alpha_i \rangle}. \quad (8)$$

Therefore, we obtain the quantization rule

$$\theta = \frac{2\pi N}{n}, \quad (n \neq 0), \quad (9)$$

with N being an integer.

3. References

1. H. Zhang, *Preprint* **LBL-32491**, June 1992; **LBL-32531**, June 1992.
2. H. Zhang, *Preprint* **LBL-32595**, July 1992.
3. R. D. Peccei and H. R. Quinn, *Phys. Rev. Lett.* **38**, 1440 (1977); *Phys. Rev. D* **16**, 1791 (1977); S. Weinberg, *Phys. Rev. Lett.* **40**, 223 (1978); F. Wilczek, *Phys. Rev. Lett.* **40**, 271 (1978).
4. Y. S. Wu and A. Zee, *Nucl. Phys.* **B258**, 157 (1985).
5. R. Jackiw, in *Relativity, Groups and Topology II*, (Les Houches 1983, ed. B. S. DeWitt and R. Stora, North-Holland, Amsterdam, 1984).
6. E. Witten, *Phys. Lett.* **B86**, 283 (1979).
7. H. Zhang, *Phys. Rev.* **D36**, 1868 (1987).
8. H. Pagels, *Phys. Rev.* **D13**, 343(1976); W. Marciano and H. Pagels, *Phys. Rev.* **D14**, 531(1976).

9. P. Nelson and A. Manohar, *Phys. Rev. Lett.* **50**, 943 (1983); A. Balachandran et al, *Phys. Rev. Lett.* **50**, 1553 (1983); A. Abouelsaood, *Phys. Lett.* **B125**, 467 (1983).